

Phone: 208.343.6458 Fax: 208.343.5031 Web: IdahoGastro.com

Small Bowel (Ileoscopy Prep) Instructions

#### Read and follow all instructions carefully

Important: Do not eat breakfast the day prior to your procedure

## **PURCHASE AT THE PHARMACY**

- 1-119 gram bottle of MiraLAX (non-prescription)
- 32 ounces of a sports drink of your choice (any color except red)
- 2 tablets Zofran (prescription for nausea will be sent in closer to your procedure)

## PREPARATION FOR YOUR PROCEDURE

- **Continue all** aspirin and anti-inflammatory medications including ibuprofen, Motrin, Aleve, naproxen, Naprosyn, sulindac, piroxicam, Feldene, indomethacin, diclofenac, and Voltaren. Tylenol (acetaminophen) is also safe to continue.
- If you take a prescribed blood thinner, you will need holding instructions from our office.
- Arrange to have a driver take you home after the procedure. You will not be allowed to drive after receiving sedation.
- If you have <u>diabetes</u> or are taking medications for weight loss, use the following instructions:
  - Fasting (going without food) is required for this procedure, monitor your blood sugar and follow these directions
    - Insulin Pump: Leave on, if you have questions refer to your endocrinologist.
    - Long-acting Insulin: half dose the evening before procedure. Do not take the morning of your procedure.
    - Short-acting insulin: dose as necessary with blood sugar monitoring.
    - Oral diabetic medication: do not take the morning of your procedure.
    - GLP1: Do not take the day of your procedure (may take day before or day after).
    - If you are concerned about your blood sugar dropping, consider purchasing glucose tablets.

# CALL MON-FRI 7 AM – 5 PM OR CALL OUR AFTER HOURS NUMBER

Meridian Endoscopy Center (208) 695-2100
Boise Endoscopy Center (208) 342-7169
Canyon County Endoscopy Center (208) 954-8218
After Hours (208) 343-6458

#### **5 Days Before Your Appointment**

- No nuts or seeds.
- Stop all fiber supplements including Metamucil, Citrucel, Benefiber, FiberChoice and flax seeds.
- Stop all iron supplements, fish oil, CBD oil, or any oil-based supplements.

# **The Day Before Your Appointment**

- Step 1
  - From the moment you wake up, start a clear liquid diet. Do not eat breakfast. Avoid red beverages.
    - This includes clear juices (apple, grape, strained orange, strained lemonade), tea, coffee (no milk or creamer), soft drinks, sports drinks (Gatorade, Powerade), clear broth or bouillon, Jell-O, and Popsicles.
    - Avoid red drinks, red Jell-O, and dairy products.
    - Consume large amounts of clear fluids to increase success of the bowel flush, help avoid hunger and avoid dehydration.
- Step 2
  - 5:00 pm, mix the bottle of MiraLAX (119 g or 7 capfuls) into 64 ounces (8 Cups) of your selected sports drink. Shake until completely dissolved.
  - o To decrease the chance of nausea, you may take one Zofran tablet 30-60 minutes prior to drinking your prep.
  - o Drink one 8 ounce glass every 10-15 minutes until you have finished the 64 ounces (8 cups).
  - If no stools occur within 2 hours of completion, repeat this step. This will require you to purchase more MiraLAX and sports drink.
  - If you experience nausea while taking your prep, stop drinking and allow the nausea to settle. You may take a second
     Zofran tablet to help ease your nausea. When you resume drinking, drink slower or through a straw to help avoid nausea. If vomiting occurs, please call us at the numbers above.
  - o Continue drinking clear liquids. This helps your prep be more successful.

425 W. Bannock • Boise, Idaho 83702 2235 E. Gala • Meridian, Idaho 83642 1216 Garrity • Nampa, Idaho 83687



Phone: 208.343.6458 Fax: 208.343.5031 Web: IdahoGastro.com

# **The Day of Your Appointment**

- Stools should be watery, yellow and see-through. If you have concerns about the quality of your prep, please call us.
- Continue consuming <u>clear</u> liquids to avoid dehydration <u>up until 4 hours prior to your scheduled procedure.</u>
- 4 hours prior to your procedure STOP drinking fluid. You may take small sips of water only to take your prescribed medications this is very important. Drinking within 4 hours of your procedure may result in delay or cancellation.
- Bring a list of your current medications and correct dosages to your appointment.
- NO DRIVING —you cannot drive for the remainder of the day if you are sedated. You need to have a responsible driver to take you home after your procedure. If your driver wishes to discuss the results of your procedure with the doctor, please have them accompany you to your procedure and remain on site.
- We do not recommend Uber, Lyft, or taking a taxi these are not acceptable forms of transportation.
- You will be sedated for your procedure. Please avoid alcohol and making legal decisions for the remainder of the day after your procedure.

You are scheduled with Dr		
Report at	on	_ to:
	_ Boise Endoscopy Center, 425 West Bannock Street, Boise	
	_ Meridian Endoscopy Center, 2235 East Gala Street, Meridian	
	_ Canyon County Endoscopy Center, 1216 Garrity Boulevard, Nampa	
	_ St. Luke's Boise Regional Medical Center, 190 E. Bannock Street, Boise	
	(Main hospital entrance. Out-patient Registration)	
	_ St. Luke's Meridian Medical Center, 520 S. Eagle Road, Meridian	
	(Eagle Road entrance. Out-patient Registration)	
	_ St. Luke's Nampa Medical Center, 9850 W. St. Luke's Drive, Nampa	
	(Main hospital entrance. Out-patient Registration)	

#### **Helpful Tips:**

- Clear liquids only the day before and on the day of your exam. Do NOT drink any fluids 4 hours prior to your exam.
- You must complete your entire bowel prep. We often see poor bowel preparations because patients stop drinking.
- If you have trouble drinking the prep, try sucking on ice or a lemon wedge, brush your teeth, chew gum, or rinse your mouth.
- If you finish the whole prep and your stool is brown, thick, or not clear please call us for advice.
- If you are nauseated or vomit with the prep, stop drinking to allow your stomach to settle and take the Zofran we provide.
- You may not remember your conversation with your doctor after your procedure we STRONGLY recommend your driver stays IN our facility to speak with the doctor after your test.

Your procedure time will be approximately 90 minutes from the scheduled check-in time to discharge for procedures in the Meridian, Boise, and Canyon County Endoscopy Centers. Please coordinate with your driver to alleviate delays in discharge.

Patients' stay times for hospital procedures may be longer.

We cannot be responsible for your valuables (watches or various jewelry items). Please leave them at home.

Bring all insurance cards and a photo ID with you. Please check with your insurance company regarding prior authorization.

Fee for services is required at time of check-in. Please be prepared to make a payment.

IGA recognizes that your time is valuable, and so is ours. If you need to cancel or reschedule an appointment, please do so more than 48 hours prior to your scheduled appointment arrival time at 208-343-6458. Late cancellations and no shows may be assessed a \$200 fee.